

Evaluation of fungicide seed treatments as management tools for root rot of field peas in North Dakota

Introduction

Root rots are of particular concern for field pea production in North Dakota, the largest producer of field peas (*Pisum sativum*) in the US with a planting area exceeding 500,000 A and a production value exceeding \$100 million annually (NASS, 2009). Damage from root rot pathogens is being reported in all pea-growing areas of the state and is likely to increase as field peas become an established part of the landscape. Fungicide seed treatments have been used to manage root rot pathogens on field peas, but few trials have been done in North Dakota.

Objective

The objective of this study was to evaluate the efficacy of fungicidal seed treatments on field pea in North Dakota.

Materials and Methods

Four seed-treatment trials were established at the North Dakota State University (NDSU)-Carrington Research Extension Center (CREC) near Carrington, ND, and at an on-farm location in Newburg, ND, in 2008 and 2009. Field pea cultivar D.S. Admiral was planted in seven row plots measuring approximately 1.5 m x 7.6 m on 5 and 19 May at CREC in 2008 and 2009, respectively, and on 7 and 21 May at Newburg, respectively. Planting rate was 300,000 pure live seeds acre⁻¹ in 2008 and 330,000 pure live seeds acre⁻¹ in 2009. Previous crop was spring wheat in all trials except CREC in 2009, where the previous crop was dry pea. Although the focus of the trial was root rots and fungicides, several insecticides were included.

Trials were arranged in a randomized complete block with four replications and included one or two nontreated controls and 12 fungicide and/or insecticide seed treatments, including; trifloxystrobin + metalaxyl (Trilex, BayerCropScience, Research Triangle Park, NC), mefenoxam + fludioxonil (ApronMaxx, Syngenta Crop Protection, Greensboro, NC), mefenoxam + fludioxonil + thiamethoxam (CruiserMaxxBears, Syngenta, Research Triangle Park, NC), hymexazol (Tachigaren, Bayer CropScience, Research Triangle Park, NC), fludioxonil (Maxim, Syngenta, Research Triangle Park, NC), metalaxyl (Allegiance FI, Bayer CropScience, Research Triangle Park, NC; Acquire, BASF, Research Triangle Park, NC), thiabendazole (LSP, Bayer CropScience, Research Triangle Park, NC; Mertect, Merck, Whitehouse Station, NJ), thiophanate methyl + pyraclostrobin (Acronis, BASF Research Triangle Park, NC), thiophanate methyl (Tops 30, Bayer CropScience, Research Triangle Park, NC), Imidacloprid (Gaucho 600, Bayer CropScience, Research Triangle Park, NC), and pyraclostrobin (Stamina, BASF, Research Triangle Park, NC).

Stand establishment was evaluated at least once at approximately 14 d post emergence by counting emerged plants in center two rows. Disease incidence and severity were assessed by randomly sampling ten plants approximately 4 weeks after emergence. Incidence was recorded as percent plants with root rot symptoms and severity was recorded as percent infection (visible lesions) on the taproot. Infected roots were washed, surface sterilized, and cultured and pathogens were identified using culture morphology and DNA sequencing. Plots were harvested at CREC on 6 and 18 August in 2008 and 2009, respectively, and at Newburg on 14 and 26 August, respectively. Yield, protein, test weight were evaluated after harvest. Data were analyzed using the Statistical Analysis Software (SAS Institute, Cary, NC) PROC ANOVA software (version 9.1).

Results and Discussion

Root rot was observed in all trials but was more severe at Newburg in both years. Several *Fusarium* species (mainly, *Fusarium oxysporum*, *Fusarium redolens*, *Fusarium avenaceum*, *Fusarium acuminatum*, *Fusarium graminearum*) and *Rhizoctonia solani* (AG-4) were identified in each location year (Tables 1-4). Statistical differences between non-treated controls and chemical seed treatments were observed in multiple parameters in multiple trials (Tables 1-4). A yield increase over the non-treated control(s) were observed in CREC in 2008 and 2009, and although statistical differences were not observed, yield of non-treated controls at Newburg in 2008 accounted for two of the lowest three values. Although fungicide seed treatments resulted in a yield increase over the non-treated control, two fungicide/insecticide combination treatments yielded significantly higher than fungicide only treatments in CREC 2009. Insect damage was not evaluated.

Conclusions

- ✓Yield increases in yield in two of four trials indicate that seed treatments may be an economically viable option for management of root rot.
- ✓Several *Fusarium* species and *Rhizoctonia solani* contributed to root rots in all trials.
- ✓Emphasis should be place on evaluating the efficacy of seed treatments effective against *Fusarium* species.
- ✓Insecticide seed treatments should be evaluated.

Treatment Name	Product/AI Rate	Newberg 2008								Carrington 2008							
		Plant Stand plants ft ⁻²	Root Rot Incidence	Root Rot Severity (% root infection)	Yield (bu/A)	Test Weight lbs/bu	Seed Protein %	Pathogens isolated		Plant Stand plants ft ⁻²	Root Rot Incidence	Root Rot Severity (% root infection)	Yield (bu/A)	Test weight (lbs/bu)	Seed Protein (%)	Pathogens isolated	
								<i>Fusarium</i> spp.	<i>Rhizoctonia</i> spp.							<i>Fusarium</i> spp.	<i>Rhizoctonia</i> spp.
Nontreated		8.0	30.0	9.1	28.2	62.9	23.5	5	0	7.7	12.5	0.6	50.7 a	64.8	23.4	1	0
Tops 30	0.5 fl oz / 100 lbs seed	7.8	35.0	7.2	31.8	63.0	23.4	4	0	8.8	17.5	2.0	60.7 b	64.5	23.9	4	1
Trilex Conc.	1.0 fl oz / 100 lbs seed	6.8	35.0	3.3	31.4	63.3	23.4	2	0	8.2	12.5	2.3	60.7 b	64.9	23.5	3	0
Trilex Conc. + Tops 30	1.0 + 0.5 fl oz / 100 lbs seed	7.7	30.0	5.2	32.2	63.4	23.5	3	0	8.7	10.0	0.8	63.6 c	64.9	23.5	2	0
Trilex Conc.+Gaucho 600	1.0 + 1.6 fl oz / 100 lbs seed	6.6	12.5	5.0	28.5	62.9	23.5	4	0	7.7	7.5	1.0	62.5 b	64.8	23.6	2	0
Apron Maxx	5.0 fl oz / 100 lbs seed	7.6	27.5	4.7	29.9	63.0	23.4	4	1	8.6	22.5	1.8	61.6 b	64.8	23.9	4	0
Cruiser Maxx	1.5 fl oz / 100 lbs seed	7.1	22.5	5.0	30.7	63.3	23.6	4	0	8.5	12.5	1.5	60.2 b	64.7	23.7	2	1
Mertect	3.0 fl oz / 100 lbs seed	7.1	17.5	4.0	31.8	63.5	23.5	3	0	7.6	27.5	2.8	57.9 b	64.9	23.5	2	1
Acquire + Acronis	0.3 + 1.5 oz / 100 lbs seed	7.4	15.0	8.5	32.4	62.9	23.5	4	0	9.0	2.5	0.3	60.8 b	64.2	23.5	1	0
Tachigaren	15 grams / kg	7.4	20.0	6.5	33.2	62.9	23.5	3	0	7.7	5.0	0.3	60.5 b	64.7	23.8	0	0
Maxim (4FS)	0.16 fl oz / 100 lbs seed	7.1	12.5	5.0	33.7	62.9	23.6	3	0	8.2	0.0	0.0	61.5 b	64.8	23.8	0	0
Allegiance FI	0.75 fl oz / 100 lbs seed	6.5	10.0	2.0	29.7	63.3	23.4	2	0	8.6	10.0	1.0	60 b	64.6	23.7	2	0
Apron Maxx + LSP	5.0 fl oz + 3.28 fl oz / 100 lbs	6.3	12.5	3.0	31.6	63.1	23.6	2	0	8.6	10.0	1.0	63.2 b	64.6	23.7	3	0
Nontreated		7.9	25.0	13.1	29.1	63.2	23.5	4	0	7.4	12.5	1.3	49.9 a	64.8	23.4	4	0
LSD(P=0.05)		NS	22.02	7.57	NS	NS	0.3			NS	14.81	1.67	5.5	NS	0.3		
CV		14.3	70.66	90.93	8.8	0.6	0.9			11.8	89.23	99.63	6.5	0.6	0.9		

Treatment Name	Product/AI Rate	Newberg 2009											Carrington 2009								
		Plant Stand plants ft ⁻²	Root Rot Incidence	Root Rot Severity (% root infection)	Yield (bu/A)	Test Weight lbs/bu	Seed Protein %	Root Rot Pathogens / treatment			Plant Stand plants ft ⁻²	Root Rot Incidence	Root Rot Severity (% root infection)	Yield (bu/A)	Test Weight lbs/bu	Seed Protein %	Root Rot Pathogens / treatment				
								<i>Fusarium</i> spp.	<i>Rhizoctonia</i> spp.	Unknown							<i>Fusarium</i> spp.	<i>Rhizoctonia</i> spp.	Unknown		
Nontreated	NA	5.9	67.5	31.4	63.3	65.4	21.0	8	1	3	6.2	77.5	32.4	31.1	63.8	19.9	13	0	2		
Cruiser Maxx Beans	3.0 oz / 100 lbs seed	6.6	57.5	22.8	66.3	65.0	21.1	5	0	3	6.3	62.5	22.4	53.9	64.2	18.9	10	0	0		
Apron Maxx RTA	5.0 oz / 100 lbs seed	6.3	52.5	25.5	62.6	65.7	21.2	14	1	1	5.9	50.0	17.3	40.0	64.4	19.2	12	0	0		
Apron Maxx RTA + Mertec	5.0 + 3.0 fl oz / 100 lbs seed	6.3	54.4	22.1	64.2	65.4	21.3	9	1	1	6.1	50.0	22.9	38.2	63.2	18.9	3	0	0		
Trilex 2000	1.0 fl oz / 100 lbs seed	6.1	47.5	13.6	62.2	65.5	21.2	6	1	3	5.7	55.0	19.9	41.2	64.0	19.1	4	0	0		
Trilex 2000 + Gaucho	1.0 + 1.6 oz / 100 lbs seed	6.6	52.5	27.1	66.4	65.6	21.5	9	1	1	6.6	55.0	18.0	64.3	64.1	19.4	4	0	1		
Trilex 2000 + Tops 30	1.0 + 0.5 oz / 100 lbs seed	5.7	40.0	10.5	63.1	65.6	21.3	10	0	0	6.2	50.0	18.3	36.5	63.7	19.4	4	0	0		
Acquire + Acronis	0.3 + 0.4 oz / 100 lbs seed	5.7	42.5	13.0	62.6	65.8	21.3	7	0	1	6.3	42.5	16.8	34.8	63.9	19.0	1	0	0		
Acquire + Stamina	0.3 + 1.5 oz / 100 lbs seed	5.5	47.5	16.0	60.8	65.6	21.0	6	1	3	6.5	35.0	9.4	37.0	63.9	19.0	5	0	2		
Maxim (4FS)	0.16 fl oz / 100 lbs seed	6.1	45.0	18.8	60.6	65.6	21.1	2	1	1	5.3	57.5	27.4	38.3	64.1	19.4	5	0	0		
Allegiance FI	0.75 fl oz / 100 lbs seed	6.6	52.5	24.7	61.5	65.7	21.3	10	1	1	6.3	55.0	21.0	41.8	63.0	19.3	2	0	0		
Apron XL	0.64 fl oz / 100 lbs seed	6.0	55.0	19.1	61.0	65.6	21.3	9	0	1	5.8	55.0	16.0	36.7	63.4	19.1	6	0	0		
Apron Maxx RFC + Maxim 4FS	1.5 fl oz + 0.08 fl oz / 100 lbs	6.2	60.0	27.6	62.4	65.7	21.2	13	1	0	6.5	50.0	11.7	37.6	63.8	19.2	6	0	0		
LSD(P=0.05)		0.7	16.30	12.18	NS	NS	NS				NS	16.63	10.85	7.7	NS	0.8					
CV		8.2	22.27	41.87	6.3	0.8	1.4				11.2	21.73	39.46	13.5	1.1	2.8					

Acknowledgments

The authors would like to thank Tim Indergaard and other personnel at the Carrington Research Extension Center for technical and logistic support, the Pulse-Crop Research and Extension Groups in the Plant Pathology Department at NDSU for technical and logistical support, Dr. Mary Burrows at Montana State University for technical support, Cliff and Tom Issendorf for on-farm collaboration, and agro-chemical companies for fungicides and insecticides used in this trial. This research was funded by Northern Pulse Growers Association and the USDA Cool Season Food Legume program.